



## New Study: For Stable Heart Disease, Lifestyle Changes and Meds Just as Effective as Bypass, Stents

By Courtney Davison  
November 25 2019

For people with stable coronary artery disease, stents and bypass surgery are no more effective at reducing rates of heart attack and death than lifestyle changes and medication alone, a new large-scale federally funded study shows.

The [ISCHEMIA trial](#) examined the efficacy of surgical intervention in patients with ischemic heart disease, also known as coronary artery disease. IHD occurs when plaque accumulates in the heart's major blood vessels, reducing its blood flow. It is the most common type of heart disease in the United States and the leading cause of death worldwide, resulting in more than 9 million deaths annually.

A collaboration between researchers at NYU Langone Health and 15 other institutions, the ISCHEMIA study followed 5,000 subjects with moderate to severe IHD. Participants were split into two groups: One group received only lifestyle counseling and medical therapy. The other group received [stents](#) or bypass surgery in addition to lifestyle counseling and medical therapy. After a median duration of four and a half years, researchers found no significant difference in rates of heart attack and death between the two groups.

Monica Aggarwal, MD, director of [Integrative Cardiology and Prevention at the University of Florida](#), says she was not surprised by the results. "Many of us [cardiologists] who practice more conservatively believe that lifestyle modification and medications are better than [surgical] intervention, and this trial gave us validated information."

The ISCHEMIA authors [presented](#) their research at the annual American Heart Association conference in November 2019, noting that these findings do not apply to people having a heart attack or those with severe chest pain—cases in which emergency surgery may be indicated.

This is not the first study to conclude that surgical intervention should be used more sparingly for patients with stable cardiovascular disease: A 2017 study [found stents were ineffective for chest pain](#).

Aggarwal notes that a lot of people mistakenly view surgical interventions as "no big deal." "These are not benign procedures, by any means." Aggarwal adds that while stents and catheterizations are relatively low-risk, any time a surgeon goes into the arteries, there is the potential for trauma or damage to the arteries. "Many people are getting these interventions that don't need them, and that is what this study showed," says Aggarwal.

For patients with stable ischemic heart disease, Aggarwal advises a combined approach of medical therapy and healthy lifestyle initiatives, including a [whole-food, plant-based diet](#). "We have to remember that medications and lifestyle changes are complementary," Aggarwal says.

"I have seen patients with stable ischemic heart disease whom I have treated with a whole-food, plant-based diet make marked improvements in symptoms and activity," says Columbus Batiste, MD, chief of cardiology at Kaiser Permanente Riverside. "These clinical improvements have allowed them to forego stents and surgeries and also [to reduce] medications."

A [study](#) presented at the 2017 AHA conference found that a plant-based diet was associated with a 42 percent reduction in the risk of heart failure among people without previously diagnosed heart disease or heart failure. A [2014 study](#) of 198 patients with cardiovascular disease found that subjects who adopted a plant-based diet had a much lower rate of subsequent cardiac events compared with the subjects who did not adopt a plant-based diet.